



**COLON, STOMACH & LIVER CENTER, LLC | LOUDOUN ENDOSCOPY GROUP**

Satinder Gill, M.D. | Hiwot Desta, M.D. | Rahul Kataria, M.D.

Colleen Nappi, CFNP | Ndeye Kone PA-C

19455 Deerfield Ave., Suite 201 | Lansdowne, VA 20176

(P) 703.723.3670 | (F) 877.325.2018



**Preparation for Colonoscopy**

**\*PLEASE READ THOROUGHLY\***

**7 Days Prior to Procedure**

- Stop taking oral iron, fish oil, and aspirin medication.
- Please monitor for multivitamins that may contain iron.
- **AVOID** nuts, seeds, and popcorn.
- If you take medication to thin your blood, make sure you have consulted with your provider and/or cardiologist for a specific time to stop medication prior to procedure.

**1 Day Prior to Procedure**

- **CLEAR LIQUID DIET** (breakfast, lunch, and dinner) **ALL DAY**.  
 AVOID all **RED**, **PURPLE**, and **BLUE** colored drinks or liquids/jellos. Clear liquid diet includes:
 

✓ Clear broth soup	✓ Gatorade	✓ Apple or white grape juice
✓ Popsicles	✓ Power-Aid	✓ Crystal Light
✓ Jell-O (Gelatin)	✓ Tea	✓ Clear soft drinks (Sprite, club soda, ginger-ale)
✓ Italian Ices	✓ Coffee (w/o cream)	
- Take medications as directed by physician.
- Throughout the day, drink at least 8 tall glasses of clear liquids prior to beginning prep kit in the evening. Hydration is very important.
- Begin bowel prep as directed by prep solution instructions given in the office. **DO NOT FOLLOW the directions in and on the prep kit box.** Timing may vary according to which prep you are given.
- Is the prep working?
  - Bowel movements should be changing in color towards yellow and liquid (almost like urine).
- After midnight (12:00 AM), DO NOT eat or drink.

**Day of Procedure**

- In the morning, you must take important medications at least 4 hours prior to your procedure (blood pressure, post-transplant, seizure, heart rhythm, etc.) with a few small sips of water. (Unless directed otherwise)
- Do you have a driver? Ensure someone will drive you home. **No Taxi/Uber/Lyft, etc.**
- Remember to check in at the facility at the appropriate time given.
- You may eat a regular meal **AFTER** the procedure.

**KEYS** to a successful preparation:

- Staying hydrated to reduce risk of dehydration
- Eating a low fiber diet for a few days prior to prep day (avoid fiber supplements)
- If you are constipated the week of the procedure, take one packet of Miralax the evening prior to prep day.

TIP: Use a straw for drinking the solution.

What should your bowel movements look like after the prep?

- No stool present in the bowel movement
- Urine colored liquid (will NOT be clear like water)